

Physiological Response to Negative Emotions in Children with Anxiety Symptoms: Many Steps Still to be Taken

Simona Scaini,^{1,2,*} Anna Ogliari,^{2,3} and Chiara Brombin⁴

¹Department of Psychology, Sigmund Freud University, Porta di Ripa Ticinese 77, Milan, Italy

²Developmental Psychopathology Unit, Vita-Salute San Raffaele University, via Stamira d'Ancona 20, Milan, Italy

³Department of Clinical Neuroscience, San Raffaele Scientific Institute, via Stamira d'Ancona 20, Milan, Italy

⁴University Centre of Statistics in the Biomedical Sciences (CUSBS), Vita-Salute San Raffaele University, Milan, Italy

*Corresponding author: Simona Scaini, PhD, Tenured Lecturer in Developmental Psychology and Psychopathology, Department of Psychology, Sigmund Freud University, Ripa di Porta Ticinese, 7720143, Milan, Italy. Tel: +39-283241854, E-mail: s.scaini@milano-sfu.it

Received 2016 July 10; Accepted 2016 August 21.

Keywords: Anxiety, Children, Physiological Response, Emotion

Dear Editor,

Anxiety symptoms are very frequent in childhood and are associated with significant impairments in social and academic functioning (1). Thus, understanding the underlying etiological processes represents an important research task in the field (2-4). Evidence show that emotion dysregulation is a fundamental feature of anxiety disorders (5). Despite the large amount of literature available on adults, only few studies focused on the psychophysiological response induced by emotion triggering stimuli in children with anxiety (6). Physiological models hypothesized that anxiety disorders are characterized by hyperarousal in response to threatening stimuli (7). While several studies focused on endocrine response reporting an association between cortisol levels and symptoms of anxiety disorders (8, 9), little work is done to identify differential patterns of physiological response to negative, anger and fear-eliciting, and affective stimuli.

Previous studies found increased heart rate (10), adrenergic activity (11), and increased baseline zygomatic muscle tension (12) in children with anxiety. In addition, significant differences in pulse rate and systolic blood pressure during a social-evaluative task were found by Beidel et al. (13) in a community sample of children with anxiety compared to the children without anxiety. More recently, a study by Weems et al. (14) examined the physiological responses (heart rate and skin conductance) induced by a phobic/fear-eliciting stimulus; i.e., a video of a large dog, in children. Data indicated a high association between psychophysiological parameters, monitored during the task, and youth, but not parents', reports of anxiety symptoms. Thus, based on the data available in the literature, it seems that children affected by anxiety symptoms are characterized by different autonomic activity in response to threatening stimuli compared to con-

trols. However, these conclusions are derived from studies using small sample sizes. In addition, no study accounted for different types of anxiety disorders. This calls for new studies examining psychophysiological responses to negative emotion-provoking stimuli in children with anxiety. Several benefits could be derived from this research area. First of all, by jointly evaluating subjective psychometric data and objective physiological responses, a more reliable clinical tool could be developed to assess patients' symptomatology. Moreover, changes in physiological responses to emotionally charged stimuli could be used as a primary measure to evaluate treatment efficacy.

Acknowledgments

The current study was supported by the FIRB Project RBFR12VHR7 entitled "Interpreting Emotions: A Computational Tool Integrating Facial Expressions and Biosignals Based on Shape Analysis and Bayesian Networks" (Dr. Brombin recipient) and by the NARSAD Young Investigator Grant 23553 entitled "Integrating Multimodal Affective Information for Emotion Recognition in Children with Symptoms of Anxiety Disorders" (Dr. Scaini recipient).

Footnote

Disclosure of Interest: Authors declared no conflict of interest.

References

1. Lyneham HJ, Rapee RM. Evaluation and treatment of anxiety disorders in the general pediatric population: a clinician's guide. *Child Adolesc Psychiatr Clin N Am.* 2005;14(4):845-61. doi: 10.1016/j.chc.2005.05.002. [PubMed: 16171705] x.

2. Scaini S, Ogliari A, Eley TC, Zavos HM, Battaglia M. Genetic and environmental contributions to separation anxiety: a meta-analytic approach to twin data. *Depress Anxiety*. 2012;**29**(9):754–61. doi: [10.1002/da.21941](https://doi.org/10.1002/da.21941). [PubMed: [22496044](https://pubmed.ncbi.nlm.nih.gov/22496044/)].
3. Scaini S, Belotti R, Ogliari A. Genetic and environmental contributions to social anxiety across different ages: a meta-analytic approach to twin data. *J Anxiety Disord*. 2014a;**28**(7):650–6. doi: [10.1016/j.janxdis.2014.07.002](https://doi.org/10.1016/j.janxdis.2014.07.002). [PubMed: [25118017](https://pubmed.ncbi.nlm.nih.gov/25118017/)].
4. Scaini S, Pezzica E, Belotti R, Colonna S, Ogliari A. The Nature of Covariation between Separation Anxiety Symptoms and Obsessive Compulsive Symptoms in Developmental Age. *Ann Depress Anxiety*. 2014b;**1**(4):1017.
5. Hannesdottir DK, Ollendick TH. The role of emotion regulation in the treatment of child anxiety disorders. *Clin Child Fam Psychol Rev*. 2007;**10**(3):275–93. doi: [10.1007/s10567-007-0024-6](https://doi.org/10.1007/s10567-007-0024-6). [PubMed: [17705098](https://pubmed.ncbi.nlm.nih.gov/17705098/)].
6. Beidel DC. Assessing anxious emotion: A review of psychophysiological assessment in children. *Clin Psychol Rev*. 1989;**9**(6):717–36. doi: [10.1016/0272-7358\(89\)90019-6](https://doi.org/10.1016/0272-7358(89)90019-6).
7. Clark LA, Watson D. Tripartite model of anxiety and depression: psychometric evidence and taxonomic implications. *J Abnorm Psychol*. 1991;**100**(3):316–36. [PubMed: [1918611](https://pubmed.ncbi.nlm.nih.gov/1918611/)].
8. Battaglia M, Bajo S, Strambi LF, Brambilla F, Castronovo C, Vanni G, et al. Physiological and behavioral responses to minor stressors in offspring of patients with panic disorder. *J Psychiatr Res*. 1997;**31**(3):365–76. [PubMed: [9306294](https://pubmed.ncbi.nlm.nih.gov/9306294/)].
9. Gunnar MR, Vazquez DM. Low cortisol and a flattening of expected daytime rhythm: potential indices of risk in human development. *Dev Psychopathol*. 2001;**13**(3):515–38. [PubMed: [11523846](https://pubmed.ncbi.nlm.nih.gov/11523846/)].
10. Beidel DC. Psychophysiological assessment of anxious emotional states in children. *J Abnorm Psychol*. 1988;**97**(1):80–2. [PubMed: [3351116](https://pubmed.ncbi.nlm.nih.gov/3351116/)].
11. Rogeness GA, Cepeda C, Macedo CA, Fisher C, Harris WR. Differences in heart rate and blood pressure in children with conduct disorder, major depression, and separation anxiety. *Psychiatr Res*. 1990;**33**(2):199–206. doi: [10.1016/0165-1781\(90\)90074-f](https://doi.org/10.1016/0165-1781(90)90074-f).
12. Turner SM, Beidel DC, Epstein LH. Vulnerability and risk for anxiety disorders. *J Anxiety Dis*. 1991;**5**(2):151–66. doi: [10.1016/0887-6185\(91\)90026-p](https://doi.org/10.1016/0887-6185(91)90026-p).
13. Beidel DC. Determining the reliability of psychophysiological assessment in childhood anxiety. *J Anxiety Dis*. 1991;**5**(2):139–50. doi: [10.1016/0887-6185\(91\)90025-o](https://doi.org/10.1016/0887-6185(91)90025-o).
14. Weems CF, Zakem AH, Costa NM, Cannon MF, Watts SE. Physiological response and childhood anxiety: association with symptoms of anxiety disorders and cognitive bias. *J Clin Child Adolesc Psychol*. 2005;**34**(4):712–23. doi: [10.1207/s15374424jccp3404_13](https://doi.org/10.1207/s15374424jccp3404_13). [PubMed: [16232068](https://pubmed.ncbi.nlm.nih.gov/16232068/)].